Injection assessment module

Health Post Unique ID:

In.0	Child's UNIQUE ID (Composed of the 2 digit region, 2		
	digit zonal, 2 digit woreda code, 2 digit PHCU code, one digit health post one digit child's id)	Region / Zone / Woreda / PHCU / Health Post	In.0

Material required for IM injection includes:

- 1. "Injection trainer model"
- 2. "Injection box"
 - o IM medication / Saline ampoule (2 or more ml)
 - o 1 ml or 2 ml syringe
 - antiseptic swab
 - o Cotton wool swab
- 3. "Wrist watch"

no		Categories	Skip pattern	response	No
In.1	Are you trained to give IM antibiotic/ gentamycin injections in the thigh to neonate less than 1 month old?	1= Yes 2= No	Yes =skip in.3		In.1
In.2	Are you trained to give IM vaccination 1= Yes injections in the thigh to neonate less than 1 month old?		N0= skip the module		In.2
	 Instructions: Ask the assistant (e.g. community mobilizer) to sit on the bench/ chair Strap the "injection trainer" on the left thigh of the assistant Place the opened injection box in front of the health worker on the table Instructions for the health worker: Assume this is a thigh of a new-born of less then I month (pointing towards the "injection model) with all its flash, blood, and skin sensitivity. S/he is here for a IM antibiotic/ gentamycin injection (if the health worker is only trained on childhood immunization then mention "vaccine injection"). Can you please perform IM injections this newborn Any material needed for the injection is lying on the table Please feel free to ask any questions now, otherwise please proceed with the procedure 				

In.3	Hand hygiene: before		In.3
	(performs or verbally mentions)		
In.3a	Water only	1= Yes	In.3a
		2= No	
In.3b	Water and soap	1= Yes	In.3b
		2= No	
In.3c	Use of disinfectant	1= Yes	In.3c
		2= No	
In.4	Date of expiry on the ampule has been	1= Yes	In.4
	checked.	2= No	
In.5	Open the first ampule successfully.	1= Yes	In.5
		2= No	
In.6	Fill up the syringe 1 ml (The volume of the	1= Yes	In.6
	IM injection should not be more than 1ml.	2= No	
In.7	Injection site is cleaned using an alcohol	1= Yes	In.7
	swab (Skin must be swabbed at least 30	2= No	
	seconds)		
In.8	Allow the alcohol to air dry completely	1= Yes	In.8
	(At least 30 seconds).	2= No	
In.9	Selection the correct injection site	1= Yes	In.9
	(anterolateral thigh muscles - the junction	2= No	
	of the upper and middle thirds of the		
1. 10	bulkiest/ lateral part of the thigh muscle)	4 V:	1.40
In.10	Stretch the skin - secure the thigh between	1= Yes	In.10
In 11	your thumb and forefinger.	2= No 1= Yes	In 11
In.11	Pierce the skin at an angle of 90 degrees to the skin.	1= Yes 2= No	In.11
In.12	Perform aspiration to ensure there is no	1= Yes	In.12
111.12	large blood vessels in the injection sites	1= Yes 2= No	111.12
In.13	Slowly inject the medication for even	1= Yes	In.13
111.13	distribution and to minimise the infant's	2= No	111.13
	discomfort (at least 5 seconds).	2-110	
In.14	Remove the needle. Apply cotton wool ball	1= Yes	In.14
	to the injection site.	2= No	
In.15	Dispose of the needles and syringe in	1= Yes	In.15
	appropriate container	2= No	
	app. spriace container		